arXiv:1203.6813v1 [astro-ph.GA] 30 Mar 2012

Using deuterated H_3^+ and other molecular species to understand the formation of stars and planets

F.F.S. van der Tak

SRON Netherlands Institute for Space Research, Landleven 12, 9747 AD Groningen, The Netherlands

The H_3^+ ion plays a key role in the chemistry of dense interstellar gas clouds where star and planets are forming. The low temperatures and high extinctions of such clouds mak direct observations of H_3^+ impossible, but lead to large abundances of H_2D^+ and D_2H^+ which are very useful probes of the early stages of star and planet formation. Maps o H_2D^+ and D_2H^+ pure rotational line emission toward star-forming regions show tha the strong deuteration of H_3^+ is the result of near-complete molecular depletion of CNO-bearing molecules onto grain surfaces, which quickly disappears as cores warm up afte stars have formed.

In the warmer parts of interstellar gas clouds, H_3^+ transfers its proton to other neutral such as CO and N_2, leading to a rich ionic chemistry. The abundances of such species ar useful tracers of physical conditions such as the radiation field and the electron fraction Recent observations of HF line emission toward the Orion Bar imply a high electron fraction, and we suggest that observations of OH^+ and H_2O^+ emission may be used t probe the electron density in the nuclei of external galaxies.

Key words: astrochemistry – interstellar medium – stars: formation.

1. Introduction

The H_3^+ ion plays a central role in the physics and chemistry of interstella gas clouds where stars and planets are forming (Watson 1973). In cold parts o such clouds, where temperatures are $\lesssim 20$ K, H_3^+ is converted into its deuterate forms, which become the dominant forms of H_3^+ at $T \lesssim 10$ K. Here, H_3^+ can b studied directly by rotational spectroscopy of H_2D^+ and D_2H^+ in the far-infrare and submillimeter. In warm interstellar clouds ($T \sim$30-100 K), a direct study o H_3^+ is possible through vibrational spectroscopy in the mid-infrared, and thi technique is widely used for lines of sight through the diffuse interstellar mediu (see contributions by McCall, Indriolo and Geballe to this volume). To peer int highly opaque star-forming regions and to map their spatial structure, emissio from indirect probes such as HCO^+ and N_2H^+ is used, which form by proto transfer of H_3^+ to CO and N_2, respectively.

The goal of this paper is to review recent work on the astrophysical use of H_3^+ and its deuterated forms. Since the role of H_3^+ in the warm and cold regimes i so different, this review discusses them separately: § 2 covers the study of H_2D^+

and D_2H^+ in cold clouds, while §§ 3 and 4 treat probes of H_3^+ in warm regions. Space limitations prevent a complete coverage of all work since the previous H_3^+ conference in 2006; therefore the content of this review is necessarily biased toward the interests of its author. In particular, this paper does not cover the recent results on the ortho/para ratios of H_2 and H_2D^+, which may constrain the ages of pre-stellar cores; see the contribution by Pagani to this volume. This work is also of interest as it marks the first use of the detailed state-to-state rate coefficients for de-excitation of H_2D^+ in collisions with H_2 (Hugo et al. 2009). Collisions between H_3^+ and H_2 may lead to reaction (fully elastic case), excitation (fully inelastic case), or both. Thus, the rates by Hugo et al. are a major step forward for the interpretation of interstellar H_2D^+ lines, compared to the scaled radiative rates used previously (see discussion by Van der Tak et al. 2005). The next step up from Hugo's statistical approach would be full quantum scattering calculations, which would be a large effort because of the reactive nature of deuterated H_3^+.

Another important topic that this paper does not discuss in detail are observations of H_2D^+ in protoplanetary disks. The conditions in the midplanes of such disks are somewhat similar to those in pre-stellar cores: the densities are high enough ($\gtrsim 10^8\,\mathrm{cm}^{-3}$) that ultraviolet radiation does not penetrate, leading to a low temperature ($<10\,\mathrm{K}$) and a low ionization fraction. The theory of chemistry in the limit of complete depletion of CNO-bearing species (Walmsley et al. 2004) was originally developed to describe pre-stellar cores, but may be even more applicable to protoplanetary disk midplanes. Here, all but the lightest molecular species freeze out onto grain surfaces, and H_2D^+ is the only available ion to probe the ionization rate of the gas, which controls the efficiency of interaction with the magnetic field. Since the magneto-rotational instability is thought to play a major role in the dynamics of such disks, several teams have searched for H_2D^+ line emission. After an initial claimed detection by Ceccarelli et al. (2004), only upper limits have been reported by Guilloteau et al. (2006), Qi et al. (2008), Chapillon et al. (2011) and Öberg et al. (2011). See these papers for the quantitative implications of the limits, and the perspectives with future telescopes.

2. Cold gas: Spatial distribution of H_2D^+ and D_2H^+

As reviewed previously (Van der Tak 2006), the astrophysical use of deuterated H_3^+ is based on the fact that the formation of H_2D^+ and D_2H^+ from H_3^+ + HD is strongly enhanced at low temperatures ($\lesssim 20\,\mathrm{K}$). As a result, the abundance of deuterated H_3^+ peaks in the dense, cold concentrations of gas within clouds known as cloud cores. The temperature dependence is enhanced by a second effect: the main destroyer of deuterated H_3^+ is CO, which freezes out onto grain surfaces under the same dense and cold conditions. These two effect make H_2D^+ and D_2H^+ ideal tracers of the initial conditions of star formation; see Sipilä et al. (2010) for recent models of this situation. At higher temperatures ($\sim$20–30 K), reactions with CH_2D^+ and C_2HD^+ rather than H_2D^+ act to enrich molecular D/H ratios (Parise et al. 2009; Sakai et al. 2009; Öberg et al. 2012).

The importance of deuterated H_3^+ has been realized for a long time e.g., Pagani et al. 1992 but whether this potential could be used remained unclear until the first detection spectra of interstellar H_2D^+ (Stark et al. 1999; Caselli et al.

2003). Motivated by these results, Caselli et al. (2008) present the first survey of H_2D^+ line emission, toward a sample of 10 starless cores and 6 protostellar cores. Using the Caltech Submillimeter Observatory, the H_2D^+ line at 372 GHz is detected in 7 starless cores and in 4 protostellar cores. The column density of o-H_2D^+, derived from the 372 GHz line intensity, is also higher in starless cores than in protostellar cores. The brightest H_2D^+ lines are detected toward the densest and most centrally concentrated starless cores, where the CO depletion factor (measured by the ratio of CO line emission to dust continuum emission) and the deuterium fractionation (measured as the N_2D^+/N_2H^+ line ratio) are also largest

The results of this survey are consistent with the expectation that H_2D^+ traces very dense and cold gas, partly because the reaction of H_3^+ + HD proceeds mainly in the forward direction, and partly because the main destroyer of H_2D^+,CO freezes out onto grain surfaces. In particular, at gas temperatures above 15 K, low CO depletion factors and large abundance of negatively charged small dust grains or PAHs drastically reduce the deuterium fractionation to values inconsistent with those observed toward pre-stellar and protostellar cores.

Comparing their results to a chemical model, Caselli et al. (2008) find that several parameters, including the cosmic-ray ionization rate, the depletion of CO (and generally of neutral species), the volume density, the dust grain size distribution and the fraction of PAHs all affect the H_2D^+ abundance. In addition the ortho-to-para ratio of H_2 is a key parameter (Pagani et al. 2009). The current data do not allow to disentangle these effects: observations of the p-H_2D^+ ground state line would help to break this degeneracy, as would detailed measurements of the temperatures and densities of the cores as a function of radius.

The results by Caselli et al. (2008) are corroborated by mapping of H_2D^+ line emission in the Oph B2 core (Friesen et al. 2010). The column density distributions of both H_2D^+ and N_2D^+ in this core show no correlation with the total H_2 column density. Instead, the deuterium fractionation in the Oph B2 core, again measured by the N_2D^+/N_2H^+ ratio, systematically decreases with proximity to the embedded protostars, out to distances $\gtrsim$0.04 pc.

Using the multi-pixel CHAMP+ receiver on the APEX telescope, Parise et al. (2011) have successfully detected D_2H^+ emission at 692 GHz from the H-MM1 prestellar core located in the LDN 1688 cloud. The emission is detected on several pixels of the CHAMP+ array, hence extended on a scale of at least 40″ corresponding to ~4800 AU. This is the first secure detection of interstellar D_2H^+ after the tentative detection by Vastel et al. (2004). The column densities of H_2D^+ and D_2H^+ toward this core are similar, which implies strong depletion of CO model calculations indicate that 90–99% of CO is frozen on dust grains.

Other maps of H_2D^+ emission will be published soon. The region NGC 1333 IRAS4, where H_2D^+ was first detected, is part of the Spectral Legacy Survey (SLS) at the JCMT (Plume et al. 2007), which initially covered the 330–360 GHz range but was later extended to 373 GHz. This survey provides images of molecular line emission over a 2×2′ field at 15″ resolution toward four different star-forming environments. In the case of NGC 1333 IRAS4, the H_2D^+ emission is found to have a spatial distribution unlike that of other ions such as DCO^+ or N_2H^+ or the dust continuum, which peaks near the embedded protostars. The fact that H_2D^+ emission peaks away from heat sources confirms the idea that H_2D^+ is rapidly destroyed as cores warm up (E. Koumpia et al, in preparation).

The first mapping survey of H_2D^+ toward pre-stellar cores is also underway. Using the JCMT, Di Francesco et al (in preparation) have made maps of H_2D^+ line emission toward a sample of 6 pre-stellar cores, selected from the N_2D^+/N_2H^+ study by Crapsi et al. (2005) to have high N_2D^+/N_2H^+ ratios. For each of these cores, the H_2D^+ emission is found to peak at the core center. The emission is extended over $\approx 30''$, as seen in the maps and confirmed by the similarity of H_2D^+ antenna temperatures between the CSO and JCMT telescopes.

3. Warm gas: Reactive ions in star-forming regions

Whereas large abundances of interstellar H_2D^+ and D_2H^+ require low temperatures ($\lesssim 10$ K), proton transfer from H_3^+ to other molecules also takes place in warm gas. In particular, reactions of interstellar H_3^+ with species such as CO and N_2 lead to HCO^+ and N_2H^+ which are widely observed in the interstellar medium. Stable ionic species are useful as tracers of the interaction of interstellar gas with magnetic fields (e.g., Houde et al. 2004; Schmid-Burgk et al. 2004), whereas ions which react rapidly with H_2 trace other physical conditions such as the gas density and the local radiation field. The second part of this review paper discusses a few recent results in our understanding of the physics and chemistry of reactive ions in interstellar space.

(a) *The H_nO^+ puzzle*

The HIFI instrument is a heterodyne spectrometer onboard ESA's Herschel space observatory, which provides access to the spectral ranges between 480–1250 and 1410–1910 GHz at a resolution of ≈0.1 MHz. One HIFI highlight is the discovery of interstellar OH^+ and H_2O^+ (Gerin et al. 2010; Ossenkopf et al. 2010), which are intermediate steps in the ionic formation route of interstellar H_2O. This route begins with charge transfer of H^+ or H_3^+ to O, proceeds with reactions of H_2 with O^+, OH^+ and H_2O^+, and ends with the dissociative recombination of H_3O^+ which produces H_2O. Although this route dominates the formation of H_2O in the gas phase at temperatures $\lesssim 250$ K, which is the bulk part of the neutral interstellar medium, the abundances of OH^+ and H_2O^+ were predicted to be low because these species react rapidly with H_2, which is the bulk species of dense interstellar gas clouds. In other words, the rate-limiting steps of ionic H_2O formation were thought to be the formation of O^+ and the dissociative recombination of H_3O^+.

Observations with HIFI have revealed strong absorption by interstellar OH^+ and H_2O^+ on many lines of sight in our Galaxy (Neufeld et al. 2010a), and toward nearby starburst galaxies such as M82 (Weiß et al. 2010), NGC 253 and NGC 4945. The large column densities of these reactive species imply that the hydrogen in these clouds is neither purely in atomic form, because no OH^+ and H_2O^+ would be produced, nor in purely molecular form, because all OH^+ and H_2O^+ would react into H_3O^+ and H_2O. Previous models of the structure of molecular clouds, where clouds are either diffuse and hydrogen atomic, or dense and hydrogen molecular, are inconsistent with these observations and must be rejected (e.g., Snow & McCall 2006). While PDR models are able to explain the observed amounts of OH^+ and H_2O^+ with diffuse gas where most hydrogen is



Figure 1. Spectrum of the HF J=1–0 line, observed with Herschel-HIFI toward the Orion Bar
From: Van der Tak et al (2012).

atomic (Gerin, this volume), the distribution of diffuse and dense gas must be reconsidered, as already indicated by mm-wave absorption line studies of diffuse clouds (Lucas & Liszt 1996).

A major puzzle in the study of interstellar OH^+ and H_2O^+ (which I collectively call H_nO^+) is the Herschel-SPIRE spectrum of the active galactic nucleus Mrk 231 where lines of H_nO^+ appear in emission (Van der Werf et al. 2010). The large dipole moments and the small reduced masses of H_nO^+ imply high line frequencies and large radiative decay rates, so that excitation of their rotational levels requires extremely high densities and line emission is not expected to be observable. Since my previous discussion of this topic at the SMILES conference, also organized by Prof. Oka (Van der Tak 2010), a second report of H_nO^+ emission has appeared for the ultraluminous merger Arp 220 (Rangwala et al. 2011). In this galaxy, both the OH^+ and the H_2O^+ lines show a P Cygni-type profile, which consists of blueshifted absorption and redshifted emission, and which is characteristic of a wind or an outflow.

One hint at the solution of this "H_nO^+ puzzle" may be the HF molecule. Like H_nO^+, HF appears in absorption on most lines of sight through the interstellar medium of our Galaxy (Sonnentrucker et al. 2010), which is expected from the large dipole moment and small reduced mass of the molecule. The SPIRE spectrum of Mrk 231 however shows HF in emission (Van der Werf et al. 2010) and the SPIRE spectrum of Arp 220 shows a P Cygni profile for the HF line (Rangwala et al. 2011). This behaviour is the same as for H_nO^+, which suggests (but does not imply) a connection between these species.

(*b*) *Detection of HF emission*

In dense interstellar gas clouds, the bulk of the hydrogen is in the form of H_2 which is essentially unobservable at low temperatures. The common tracer for H clouds is the CO molecule, which is abundant and chemically stable, but which photodissociates at H_2 column densities below $\approx 1.5\times10^{21}$ cm^{-2} (Visser et al. 2009). While CH may be used for the intermediate (translucent cloud) regime (Sheffer et al. 2008), a proxy for H_2 at very low column densities was missing until the Herschel mission. Since the reaction of F with H_2 leading to HF is exothermic and HF photodissociation is slow, HF is expected to be the main carrier of fluorine in the gas phase, as long as $\gtrsim 10^{-6}$ of hydrogen is in the form of H_2 (Neufeld & Wolfire 2009).

The first confirmation of these predictions came from observations with the LWS instrument onboard the ISO satellite, which measured absorption in the J=2–1 line of HF toward Sgr B2 (Neufeld et al. 1997). The Herschel-HIFI instrument gives us first access to the rotational ground state of HF, and observations of widespread absorption in this line indicate an HF abundance of $(1\text{–}2)\times10^{-8}$ in diffuse clouds, close to the interstellar (and the solar) abundance of fluorine (Sonnentrucker et al. 2010). Absorption is also observed toward dense clouds, but the implied abundance is $\sim$100× lower (Phillips et al. 2010), suggesting that depletion on grain surfaces plays a role.

The large dipole moment of HF and the high frequency of its J=1–0 line imply that radiative decay to the ground state is fast, which explains why the line usually appears in absorption on lines of sight through the interstellar medium (Neufeld et al. 2010b; Monje et al. 2011a). Only very dense environments such as the inner envelopes of late-type stars show the HF line in emission (Agúndez et al. 2011). The only detection of HF emission from the Galactic interstellar medium so far is toward the Orion Bar, and it is a surprise because the H_2 density in this region is not high enough to excite the line (Van der Tak et al. 2012). Instead, collisions with electrons appear to dominate the excitation of HF; non-thermal excitation mechanisms seem less likely, as detailed in § 4 of this paper. Line emission of HF thus appears to trace regions of molecular gas with a high electron density, which in the case of the Orion Bar is caused by strong ultraviolet irradiation by the Trapezium stars.

Excitation by electrons may also apply to active galactic nuclei where the Herschel-SPIRE spectra show HF in emission such as Mrk 231 (Van der Werf et al. 2010) or as a P Cygni profile such as Arp 220 (Rangwala et al. 2011). In contrast, the Cloverleaf quasar at $z = 2.56$ shows HF in pure absorption (Monje et al. 2011b), which may imply a lower electron density. Given the importance of electron excitation for HF, the appearance of the line in emission or absorption may be a measure of the local ionization rate by ultraviolet photons (in a starburst) or by cosmic rays (in an AGN).

The similarity of the line profiles of HF and H_nO^+ in the Herschel-SPIRE spectra of Mrk 231 and Arp 220 may not be a coincidence. Electron impact excitation may also play a role for H_nO^+ in these sources. Cross sections for the inelastic collision of the e-H_nO^+ system are urgently needed to test this hypothesis. Weak OH^+ emission is also seen toward the Orion Bar, providing further evidence for this trend (Nagy et al, in prep.).

4. Alternative mechanisms to produce HF emission

(*a*) *Infrared pumping of HF*

Radiative excitation of the HF line may occur through the v=1–0 vibration mode at 2.55 μm. This wavelength is too short for resonances with PAH features which leaves dust continuum and H_2 line emission as options. Infrared pumping is not known to play a role for other molecular species in the Orion Bar, but most of these species occur deep in the molecular cloud where near-infrared radiation does not penetrate due to extinction by dust. In contrast, HF readily forms at low extinction where both dust continuum and H_2 line emission are strong.

We have calculated the positions of the v=1–0 lines of HF from the constants of Ram et al. (1996) up to the J=10 level. Three lines are close to a line of H_2: the P(1) line at 3920.312 cm^{-1}, the R(0) line at 4000.989 cm^{-1}, and the R(1) line at 4038.962 cm^{-1}. Their respective partners are the H_2 v=2–1 Q(1) line at 3920.053 cm^{-1}, the H_2 v=1–0 Q(7) line at 4000.075 cm^{-1}, and the H_2 v=1–0 Q(6) line at 4039.507 cm^{-1}. Especially the first match is close, which is encouraging because the P(1) line leads directly to the J=1 level observed with HIFI, and the matching H_2 line is one of the strongest H_2 lines in Model 14 of Black & van Dishoeck (1987), with an intensity of ≈60% of the well-known v=1–0 S(1) line at 2.12 μm. However, the separations of these line pairs of 20, 69 and 40 km s^{-1} exceed the widths of the H_2 pure rotational lines in the Orion Bar of 4–6 km s^{-1} (Allers et al. 2005). The spatial distribution of the H_2 pure rotational lines is similar to that of the rovibrational lines (Van der Werf et al. 1996), so that their widths are probably similar too; Tielens et al. (1993) already rules out a large contribution by shocks to the H_2 rovibrational line emission from the Orion Bar.

Alternatively, the infrared pumping of HF may take place by continuum radiation from dust. Following Carroll & Goldsmith (1981), efficient pumping requires that $A_{\rm vib}\epsilon f/(e^{h\nu/kT_d}-1)$ exceeds $A_{\rm rot}$, where $A_{\rm vib}$ and $A_{\rm rot}$ are the vibrational and rotational Einstein A coefficients, ϵ is the dust emissivity (assumed to be unity in the mid-infrared), T_d is the dust temperature, and f is the geometric filling factor of the dust. Since the vibrational frequency of 3961.4 cm^{-1} is high ($h\nu/k >> T_d$), the expression simplifies to $fe^{-h\nu/kT_d} > A_{\rm rot}/A_{\rm vib}$, where the right hand side is about 1/100 using $A_{\rm vib}$=203 s^{-1} from Ram et al. (1996).

If the dust radiation field fills the entire sky as seen from the HF cloud (f=1), the lower limit for the dust temperature is 1170 K. While such warm dust could exist locally in the Orion Bar, this temperature seems unreasonably high as an average over a 7200 AU region. Even transiently heated mini-grains will rarely get so warm. Smaller filling factors exacerbate this problem: for example 50% sky coverage means a dust temperature of at least 1380 K, and 10% coverage implies T_d >2340 K. Hence continuum radiation cannot do the pumping either.

Finally, we consider the possibility that the excitation of HF in the Orion Bar is influenced by radiation from the Trapezium stars. The brightest of these stars is θ^1C Ori, which is 127″ away from the HIFI position (Van Leeuwen 2007). Assuming that both objects lie 414 pc from us (Menten et al. 2007), the stellar continuum flux at the Orion Bar is 2.6 million times stronger than at the Earth. For the 2.5 μm flux density, we adopt the K band magnitude of 4.14 (Muench et al. 2002) which corresponds to 11.5 Jy at Earth and 30.3 MJy at the Bar. Dividing this flux density by 4π gives an angle-averaged radiation field J_ν of 2.4×10^{-17} $\mathrm{erg\,s^{-1}\,cm^{-2}\,Hz^{-1}\,sr^{-1}}$. Setting this equal to the Planck function at a radiation temperature T_R, we obtain $T_R = 209\,\mathrm{K}$ at $\lambda = 2.5\,\mu\mathrm{m}$, which is only slightly larger than the intensity due to average Galactic starlight at the Sun's location. The corresponding pumping rate in the HF vibrational fundamental is of order $A_{\mathrm{vib}}/\{\exp(h\nu/kT_R) - 1\} = 1.8\times10^{-10}\,\mathrm{s^{-1}}$, which is much less than the collisional excitation rates by H_2 and electrons. We conclude that infrared pumping does not play a role for HF in the Orion Bar.

(*b*) *Formation pumping of HF*

Besides infrared pumping, an alternative non-thermal excitation mechanism of HF would be chemical pumping. In this scheme, the excitation of HF is out of thermal equilibrium because the formation and destruction rates are as fast as radiative decay. This mechanism is plausible for the Orion Bar because of the high ultraviolet radiation field, which leads to a large amount of material in the regime where H_2 is abundant but the HF destruction rate by photodissociation and reaction with C^+ is of the order of the maximum formation rate.

Assuming optically thin emission, the observed HF line flux of $7.8\,\mathrm{K\,km\,s^{-1}} = 1.5\times10^{-5}\,\mathrm{erg\,cm^{-2}\,s^{-1}\,sr^{-1}}$ requires a column density of $N_{J=1} = 9.5\times10^{11}\,\mathrm{cm^{-2}}$ in the J=1 state of HF. To obtain this by chemical pumping, the HF formation rate $R = kn(\mathrm{H_2})n(\mathrm{F})/n(\mathrm{HF})$ must fulfil $R/A_{10} \times X_{\mathrm{HF}} \times N_H \geq 9.5\times10^{11}\,\mathrm{cm^{-2}}$, where A_{10} is the Einstein A coefficient of the HF J=1–0 line. The formation of HF is dominated by the reaction of H_2 with F, the rate coefficient of which is given in Table 2 of Neufeld & Wolfire (2009) and equals $k = 4.6\times10^{-12}\,\mathrm{cm^3\,s^{-1}}$ at $T = 80\,\mathrm{K}$. In strong radiation fields, destruction of HF is dominated by photodissociation, at a rate of $\zeta_d = 1.2\times10^{-10}\,\mathrm{s^{-1}}$ in the standard Draine field, so that $\zeta_d = 6\times10^{-6}\,\mathrm{s^{-1}}$ for the Orion Bar ($\chi \sim$50,000). The HF abundance follows from balancing its formation and destruction rates ($\zeta_d = R$):

$$n(\mathrm{HF})\zeta_d = kn(\mathrm{H_2})n(\mathrm{F}) \tag{4.1}$$

where the sum of the abundances $X(\mathrm{HF}) + X(\mathrm{F})$ is constrained by the total amount of fluorine, 1.8×10^{-8} relative to H. The product $R/A_{10}\times X(\mathrm{HF})$ reaches its maximum of 4.5×10^{-12} for very high densities, leading to a maximum $N_{J=1} = 4.5\times10^{11}\,\mathrm{cm^{-2}}$ when assuming a total gas column density of $10^{23}\,\mathrm{cm^{-2}}$. For gas densities of $10^6\,\mathrm{cm^{-3}}$ this drops by a factor of two, for a gas density of $10^5\,\mathrm{cm^{-3}}$ by another factor of ten. Higher temperatures lower the required densities approximately proportionally.

The computation above assumes that every molecule formation leads to an excitation of the J=1 state. This premise is reasonable, if not intuitive. The formation of HF releases 1.36 eV in each reaction, which is well above the energy of the v=1 state, so that all rotational levels in the vibrational ground state will be about equally populated. The relative fraction of molecules formed in J=0 will be very small, and the radiative decay of all other molecules passes through the $J = 1$ state so that they contribute to the observed J=1–0 emission.

We have calculated the formation rate of HF in a self-consistent model of fluorine chemistry using the KOSMA-τ code (Röllig et al. 2006). The rate equations are solved for a spherical clump of gas and dust with a density of n_H=$10^5\,\mathrm{cm^{-3}}$ at the outer radius (A_V=0) and increasing to $1.1\times10^6\,\mathrm{cm^{-3}}$ at the



Figure 2. Chemical model of the Orion Bar as a function of A_V, using the UMIST reactio rates (dashed lines) or the H_2 + F rate from Neufeld et al. (2005) (solid lines). Top: formatio rate of HF due to reactions of F with H_2 in its v=0 and v=1 states. Middle: absolute densitie of H_2 and HF The apparent dip in the HF abundance near $A_V = 3$ when using UMIST rate is due to incorrect temperature dependence of the H_2 + F reaction rate. Bottom: calculate temperatures of gas and dust. The distribution of HF is unlike that of OH^+ and H_2O^+, whic for this χ peak at $0.05 < A_V < 1$ and disappear at $A_V > 1$ where H_2 and HF are abundant (M Röllig, priv. comm.). The observational similarity between H_nO^+ and HF is thus likely related t the non-uniform structure of the ISM and similar excitation requirements, rather than chemica similarities.

clump center, so that the total clump mass is $1\,M_\odot$. The dust model is number 7 of Weingartner & Draine (2001), equivalent to R_V=3.1, and the formation of H_2 is according to Cazaux & Tielens (2002) with formation on PAH surfaces suppressed. The total abundance of fluorine in the Orion region is taken from 6.68×10^{-9} (Simón-Díaz & Stasińska 2011), which is lower than the average value from Neufeld & Wolfire (2009) by about a factor of 3.

Figure 2 shows the results of our calculations, which consider both ground-state and vibrationally excited H_2. Collision of F with H_2 (v>0) dominates the formation of HF until A_V=0.0036, after which point ground-state H_2 takes over. The full model confirms the numbers from the estimate above with a formation rate R of about 10^{-5} s^{-1}, and approximately half of the fluorine in the form of HF in the reactive zone between $A_V \approx 0.01 \ldots 2$ containing most of the mass. Hence, we find that chemical pumping of the HF $1-0$ transition is effective in the Orion Bar, but that estimates based on current molecular data fall short by a factor of a few. This could be due to uncertainties in the total fluorine abundance, or higher temperatures and radiation fields in the interclump gas.

5. Conclusions and future work

The study of interstellar deuterated H_3^+ has come a long way in the past 16 years, and the conferences that Prof. Oka has organized every six years provide clear milestones of this progress. At the 88th birthday of the discovery of H_3^+ in 2000, the first detection of interstellar H_2D^+ was announced (Stark et al. 1999), which followed shortly after the discovery of interstellar H_3^+ itself (Geballe & Oka 1996). The 94th birthday saw the first report of strong H_2D^+ emission from cold pre-stellar cores (Caselli et al. 2003), the first H_2D^+ map of LDN 1544 by Vastel et al. (2006) and the first tentative detection of interstellar D_2H^+ (Vastel et al. 2004). This year, at the 100th birthday, we are seeing the confirmation of interstellar D_2H^+ (Parise et al. 2011), the first survey of H_2D^+ emission (Caselli et al. 2008), and mapping surveys of H_2D^+ are ongoing.

The natural question to ask now is: What will the 106th birthday in 6 years' time bring? The most likely development in the next years are high-resolution maps of pre-stellar H_2D^+ with the ALMA interferometer, which is now about half-way completed and has started its Early Science program. In fact, the tuning range of the ALMA receivers was especially extended to include the 372 GHz line of o-H_2D^+, following the CSO detection of strong emission in this line in the pre-stellar core LDN 1544. The ALMA telescope will also likely make the first detection of H_2D^+ in a protoplanetary disk, and hence constrain the ionization fraction of the disk midplane. Another possibility is a survey of the p-D_2H^+ line at 692 GHz in pre-stellar cores with ALMA, although the emission is probably quite weak in many cases. Third is the new capability of the GREAT receiver on the SOFIA airborne observatory, which gives access to the p-H_2D^+ ground state line near 1370 GHz. The science operations of SOFIA have started in 2011, and GREAT is among the first-light instruments.

The GREAT instrument may also be used for searches for the ground-state line of o-D_2H^+ at 1477 GHz. A search for this line with HIFI in the framework of the CHESS program (Ceccarelli et al. 2010) was unsuccessful (C. Vastel, priv.

comm.). Both the p-H_2D^+ 1370 GHz line and the o-D_2H^+ 1477 GHz line are unlikely to appear in emission: their upper level energies of $\sim$70 K and radiative decay rates of $\sim 3\times10^{-3}\,s^{-1}$ imply that their excitation requires higher densities and temperatures than occur in regions where the species are abundant. However it is possible that the GREAT observations will reveal absorption by p-H_2D^+ and o-D_2H^+ near 1400 GHz. Appearance in absorption requires a source of background continuum, which the warm central regions of young protostellar cores may provide, as well as sufficient abundance of H_2D^+ amd D_2H^+ in the cold outer layers of these cores. Such conditions may well occur along some lines of sight, as the example of H_2D^+ absorption at 372 GHz toward IRAS 16293 shows (Stark et al. 2004). These observations would be a new astronomical use of deuterated H_3^+ to probe physical conditions in star-forming regions.

In warm regions of the interstellar medium, hydride molecules such as HF and reactive ions such as OH^+ and H_2O^+ will continue to play an important role in characterizing the physical conditions of the gas. Mapping the spatia distribution of HF in the Orion Bar and other photon-dominated regions with Herschel-HIFI will help to clarify the contributions by atomic and molecular layers to the HF emission. The Herschel-SPIRE spectra of other galactic nuclei will help us to understand the origin of OH^+ and H_2O^+ emission by probing a range of environments. After the end of the Herschel mission in ≈March 2013, ground-based observations of OH^+ such as pioneered by Wyrowski et al. (2010) will be invaluable as probes of interstellar physics and chemistry, supported by other reactive ions that are observable from the ground such as SH^+ (Menten et al 2011), CO^+ (Stäuber & Bruderer 2009) and HOC^+ (Rizzo et al. 2003). Clearly the use of deuterated H_3^+ and other molecular ions to probe star-forming regions has a bright future!

Acknowledgment

I thank M. Röllig for input, and L. Pagani and D. Lis for comments on the manuscript.

References

Agúndez, M., Cernicharo, J., Waters, L. B. F. M., et al. 2011, A&A, 533, L6
Allers, K. N., Jaffe, D. T., Lacy, J. H., Draine, B. T., & Richter, M. J. 2005, ApJ, 630, 368
Black, J. H. & van Dishoeck, E. F. 1987, ApJ, 322, 412
Carroll, T. J. & Goldsmith, P. F. 1981, ApJ, 245, 891
Caselli, P., van der Tak, F. F. S., Ceccarelli, C., & Bacmann, A. 2003, A&A, 403, L37
Caselli, P., Vastel, C., Ceccarelli, C., et al. 2008, A&A, 492, 703
Cazaux, S. & Tielens, A. G. G. M. 2002, ApJL, 575, L29
Ceccarelli, C., Bacmann, A., Boogert, A., et al. 2010, A&A, 521, L22
Ceccarelli, C., Dominik, C., Lefloch, B., Caselli, P., & Caux, E. 2004, ApJL, 607, L51
Chapillon, E., Parise, B., Guilloteau, S., & Du, F. 2011, A&A, 533, A143
Crapsi, A., Caselli, P., Walmsley, C. M., et al. 2005, ApJ, 619, 379
Friesen, R. K., Di Francesco, J., Myers, P. C., et al. 2010, ApJ, 718, 666
Geballe, T. R. & Oka, T. 1996, Nature, 384, 334
Gerin, M., de Luca, M., Black, J., et al. 2010, A&A, 518, L110

Guilloteau, S., Piétu, V., Dutrey, A., & Guélin, M. 2006, A&A, 448, L5
Houde, M., Dowell, C. D., Hildebrand, R. H., et al. 2004, ApJ, 604, 717
Hugo, E., Asvany, O., & Schlemmer, S. 2009, J.Chem.Phys., 130, 164302
Lucas, R. & Liszt, H. 1996, A&A, 307, 237
Menten, K. M., Reid, M. J., Forbrich, J., & Brunthaler, A. 2007, A&A, 474, 515
Menten, K. M., Wyrowski, F., Belloche, A., et al. 2011, A&A, 525, A77
Monje, R. R., Emprechtinger, M., Phillips, T. G., et al. 2011a, ApJL, 734, L23
Monje, R. R., Phillips, T. G., Peng, R., et al. 2011b, ApJL, 742, L21
Muench, A. A., Lada, E. A., Lada, C. J., & Alves, J. 2002, ApJ, 573, 366
Neufeld, D. A., Goicoechea, J. R., Sonnentrucker, P., et al. 2010a, A&A, 521, L10
Neufeld, D. A., Sonnentrucker, P., Phillips, T. G., et al. 2010b, A&A, 518, L108
Neufeld, D. A. & Wolfire, M. G. 2009, ApJ, 706, 1594
Neufeld, D. A., Wolfire, M. G., & Schilke, P. 2005, ApJ, 628, 260
Neufeld, D. A., Zmuidzinas, J., Schilke, P., & Phillips, T. G. 1997, ApJL, 488, L141
Öberg, K. I., Qi, C., Wilner, D. J., & Andrews, S. M. 2011, ApJ, 743, 152
Öberg, K. I., Qi, C., Wilner, D. J., & Hogerheijde, M. R. 2012, ArXiv e-prints
Ossenkopf, V., Müller, H. S. P., Lis, D. C., et al. 2010, A&A, 518, L111
Pagani, L., Salez, M., & Wannier, P. G. 1992, A&A, 258, 479
Pagani, L., Vastel, C., Hugo, E., et al. 2009, A&A, 494, 623
Parise, B., Belloche, A., Du, F., Güsten, R., & Menten, K. M. 2011, A&A, 526, A31
Parise, B., Leurini, S., Schilke, P., et al. 2009, A&A, 508, 737
Phillips, T. G., Bergin, E. A., Lis, D. C., et al. 2010, A&A, 518, L109
Plume, R., Fuller, G. A., Helmich, F., et al. 2007, PASP, 119, 102
Qi, C., Wilner, D. J., Aikawa, Y., Blake, G. A., & Hogerheijde, M. R. 2008, ApJ, 681, 1396
Ram, R. S., Morbi, Z., Guo, B., et al. 1996, ApJS, 103, 247
Rangwala, N., Maloney, P. R., Glenn, J., et al. 2011, ApJ, 743, 94
Rizzo, J. R., Fuente, A., Rodríguez-Franco, A., & García-Burillo, S. 2003, ApJL, 597, L153
Röllig, M., Ossenkopf, V., Jeyakumar, S., Stutzki, J., & Sternberg, A. 2006, A&A, 451, 917
Sakai, N., Sakai, T., Hirota, T., & Yamamoto, S. 2009, ApJ, 702, 1025
Schmid-Burgk, J., Muders, D., Müller, H., & Brupbacher-Gatehouse, B. 2004, A&A, 419, 949
Sheffer, Y., Rogers, M., Federman, S. R., et al. 2008, ApJ, 687, 1075
Simón-Díaz, S. & Stasińska, G. 2011, A&A, 526, A48
Sipilä, O., Hugo, E., Harju, J., et al. 2010, A&A, 509, A98
Snow, T. P. & McCall, B. J. 2006, ARAA, 44, 367
Sonnentrucker, P., Neufeld, D. A., Phillips, T. G., et al. 2010, A&A, 521, L12
Stark, R., Sandell, G., Beck, S. C., et al. 2004, ApJ, 608, 341
Stark, R., van der Tak, F. F. S., & van Dishoeck, E. F. 1999, ApJL, 521, L67
Stäuber, P. & Bruderer, S. 2009, A&A, 505, 195
Tielens, A. G. G. M., Meixner, M. M., van der Werf, P. P., et al. 1993, Science, 262, 86
Van der Tak, F. 2010, ArXiv e-prints 1011.0568
Van der Tak, F. F. S. 2006, Royal Soc. of London Phil. Trans. Ser. A, 364, 3101
Van der Tak, F. F. S., Caselli, P., & Ceccarelli, C. 2005, A&A, 439, 195
Van der Tak, F. F. S., Ossenkopf, V., Nagy, Z., et al. 2012, A&A, 537, L10
Van der Werf, P. P., Isaak, K. G., Meijerink, R., et al. 2010, A&A, 518, L42
Van der Werf, P. P., Stutzki, J., Sternberg, A., & Krabbe, A. 1996, A&A, 313, 633
Van Leeuwen, F. 2007, A&A, 474, 653
Vastel, C., Caselli, P., Ceccarelli, C., et al. 2006, ApJ, 645, 1198
Vastel, C., Phillips, T. G., & Yoshida, H. 2004, ApJL, 606, L127
Visser, R., van Dishoeck, E. F., & Black, J. H. 2009, A&A, 503, 323
Walmsley, C. M., Flower, D. R., & Pineau des Forêts, G. 2004, A&A, 418, 1035
Watson, W. D. 1973, ApJL, 183, L17
Weingartner, J. C. & Draine, B. T. 2001, ApJ, 548, 296
Weiß, A., Requena-Torres, M. A., Güsten, R., et al. 2010, A&A, 521, L1

Wyrowski, F., Menten, K. M., Güsten, R., & Belloche, A. 2010, A&A, 518, A26